



## Agios Announces Publication of TIBSOVO® Phase 3 Data in *The Lancet Oncology* Demonstrating Significant Improvement in Progression-Free Survival Compared to Placebo in Previously Treated IDH1-Mutant Cholangiocarcinoma Patients

May 19, 2020

– TIBSOVO® Is the First and Only IDH1 Inhibitor to Be Evaluated in a Phase 3 Clinical Trial for IDH1-Mutant Cholangiocarcinoma –

CAMBRIDGE, Mass., May 19, 2020 (GLOBE NEWSWIRE) -- Agios Pharmaceuticals, Inc. (NASDAQ: AGIO), a leader in the field of cellular metabolism to treat cancer and rare genetic diseases, today announced that *The Lancet Oncology* has published data from its global Phase 3 ClarIDHy study of TIBSOVO® (ivosidenib) in previously treated cholangiocarcinoma patients with an isocitrate dehydrogenase 1 (IDH1) mutation. The study met its primary endpoint, demonstrating a statistically significant improvement in progression-free survival (PFS) in patients randomized to TIBSOVO® compared with placebo patients. The safety profile observed in the study was consistent with previously published data. Data from this study were [previously presented](#) at the European Society for Medical Oncology Congress (ESMO), held in September 2019 in Barcelona, Spain.

The publication can be accessed at the following link: [https://www.thelancet.com/journals/lanonc/article/PIIS1470-2045\(20\)30157-1/fulltext](https://www.thelancet.com/journals/lanonc/article/PIIS1470-2045(20)30157-1/fulltext)

"We began researching IDH inhibition as a potential therapeutic approach for solid tumors more than five years ago with the goal of innovating on behalf of patients, including those with advanced cholangiocarcinoma, who desperately need new treatment options," said Chris Bowden, M.D., chief medical officer at Agios. "We are pleased that the ClarIDHy study supports the potential utility of TIBSOVO® for this patient population and are focused on filing our supplemental new drug application for previously treated IDH1-mutant cholangiocarcinoma patients between the end of this year and mid-2021."

### ClarIDHy Phase 3 Trial

The ClarIDHy trial is a global, randomized Phase 3 trial in previously treated IDH1-mutant cholangiocarcinoma patients who have documented disease progression following one or two systemic therapies in the advanced setting. Patients were randomized 2:1 to receive either single-agent TIBSOVO® 500 mg once daily or placebo with crossover to TIBSOVO® permitted at the time of documented radiographic progression per RECIST 1.1. At the time of the primary analysis, a total of 185 patients were randomized, with 124 patients in the TIBSOVO® arm and 61 patients in the placebo arm. Thirty-five patients randomized to placebo (57.4%) crossed over to open-label TIBSOVO® upon radiographic disease progression and unblinding.

Results of the study were as follows:

- Median PFS for patients randomized to TIBSOVO® was 2.7 months compared to 1.4 months with placebo (hazard ratio [HR]=0.37; 95% CI [0.25, 0.54], one-sided  $p < 0.0001$ ) as assessed by independent radiology review.
- The estimated PFS rate was 32% at six months and 22% at 12 months for patients randomized to TIBSOVO®, while no patients randomized to placebo were free from progression beyond six months as of the data cut-off.
- The most common treatment-emergent adverse events (AEs) of any grade for the 121 patients who received TIBSOVO® were nausea (36%), diarrhea (31%) and fatigue (26%).
- Less than third of patients experienced serious AEs in either arm (30% of 121 who received TIBSOVO® versus 22% of 59 patients who received placebo).
- The most common Grade 3 or worse AE in both treatment groups was ascites (nine [7%] of 121 patients who received TIBSOVO® and four [7%] of 59 patients who received placebo).
- Patients randomized to placebo experienced a significantly greater decline in physical functioning from baseline compared to patients randomized to ivosidenib on the first day of their second 28-day treatment cycle based on European Organisation for Research and Treatment of Cancer Quality of Life Questionnaire Core 30 (EORTC QLQ-C30) Physical Functioning subscale scores ( $p=0.0059$ ).

TIBSOVO® is not approved in any country for the treatment of patients with previously treated advanced IDH1-mutant cholangiocarcinoma.

### About Cholangiocarcinoma

Cholangiocarcinoma (CC) is a rare cancer of the bile ducts within and outside of the liver. Cases that occur within the liver are known as intrahepatic cholangiocarcinoma (IHCC) and those that occur outside the liver are considered extrahepatic. Mutations in IDH1 occur in up to 20% of IHCC cases. Current treatment options for localized disease include surgery, radiation and/or other ablative treatments. There are no approved systemic therapies for IDH1-mutated cholangiocarcinoma and limited chemotherapy options are available in the advanced setting. Gemcitabine-based chemotherapy is often recommended for newly diagnosed advanced or metastatic disease.

### About TIBSOVO® (ivosidenib)

TIBSOVO® is indicated for the treatment of acute myeloid leukemia (AML) with a susceptible isocitrate dehydrogenase-1 (IDH1) mutation as detected by an FDA-approved test in:

- Adult patients with newly-diagnosed AML who are  $\geq 75$  years old or who have comorbidities that preclude use of intensive

induction chemotherapy.

- Adult patients with relapsed or refractory AML.

## IMPORTANT SAFETY INFORMATION

### WARNING: DIFFERENTIATION SYNDROME

**Patients treated with TIBSOVO® have experienced symptoms of differentiation syndrome, which can be fatal if not treated. Symptoms may include fever, dyspnea, hypoxia, pulmonary infiltrates, pleural or pericardial effusions, rapid weight gain or peripheral edema, hypotension, and hepatic, renal, or multi-organ dysfunction. If differentiation syndrome is suspected, initiate corticosteroid therapy and hemodynamic monitoring until symptom resolution.**

### WARNINGS AND PRECAUTIONS

**Differentiation Syndrome: See Boxed WARNING.** In the clinical trial, 25% (7/28) of patients with newly diagnosed AML and 19% (34/179) of patients with relapsed or refractory AML treated with TIBSOVO® experienced differentiation syndrome. Differentiation syndrome is associated with rapid proliferation and differentiation of myeloid cells and may be life-threatening or fatal if not treated. Symptoms of differentiation syndrome in patients treated with TIBSOVO® included noninfectious leukocytosis, peripheral edema, pyrexia, dyspnea, pleural effusion, hypotension, hypoxia, pulmonary edema, pneumonitis, pericardial effusion, rash, fluid overload, tumor lysis syndrome, and creatinine increased. Of the 7 patients with newly diagnosed AML who experienced differentiation syndrome, 6 (86%) patients recovered. Of the 34 patients with relapsed or refractory AML who experienced differentiation syndrome, 27 (79%) patients recovered after treatment or after dose interruption of TIBSOVO®. Differentiation syndrome occurred as early as 1 day and up to 3 months after TIBSOVO® initiation and has been observed with or without concomitant leukocytosis.

If differentiation syndrome is suspected, initiate dexamethasone 10 mg IV every 12 hours (or an equivalent dose of an alternative oral or IV corticosteroid) and hemodynamic monitoring until improvement. If concomitant noninfectious leukocytosis is observed, initiate treatment with hydroxyurea or leukapheresis, as clinically indicated. Taper corticosteroids and hydroxyurea after resolution of symptoms and administer corticosteroids for a minimum of 3 days. Symptoms of differentiation syndrome may recur with premature discontinuation of corticosteroid and/or hydroxyurea treatment. If severe signs and/or symptoms persist for more than 48 hours after initiation of corticosteroids, interrupt TIBSOVO® until signs and symptoms are no longer severe.

**QTc Interval Prolongation:** Patients treated with TIBSOVO® can develop QT (QTc) prolongation and ventricular arrhythmias. One patient developed ventricular fibrillation attributed to TIBSOVO®. Concomitant use of TIBSOVO® with drugs known to prolong the QTc interval (e.g., anti-arrhythmic medicines, fluoroquinolones, triazole anti-fungals, 5-HT<sub>3</sub> receptor antagonists) and CYP3A4 inhibitors may increase the risk of QTc interval prolongation. Conduct monitoring of electrocardiograms (ECGs) and electrolytes. In patients with congenital long QTc syndrome, congestive heart failure, or electrolyte abnormalities, or in those who are taking medications known to prolong the QTc interval, more frequent monitoring may be necessary.

Interrupt TIBSOVO® if QTc increases to greater than 480 msec and less than 500 msec. Interrupt and reduce TIBSOVO® if QTc increases to greater than 500 msec. Permanently discontinue TIBSOVO® in patients who develop QTc interval prolongation with signs or symptoms of life-threatening arrhythmia.

**Guillain-Barré Syndrome:** Guillain-Barré syndrome occurred in <1% (2/258) of patients treated with TIBSOVO® in the clinical study. Monitor patients taking TIBSOVO® for onset of new signs or symptoms of motor and/or sensory neuropathy such as unilateral or bilateral weakness, sensory alterations, paresthesias, or difficulty breathing. Permanently discontinue TIBSOVO® in patients who are diagnosed with Guillain-Barré syndrome.

### ADVERSE REACTIONS

- The most common adverse reactions including laboratory abnormalities (≥20%) were hemoglobin decreased (60%), fatigue (43%), arthralgia (39%), calcium decreased (39%), sodium decreased (39%), leukocytosis (38%), diarrhea (37%), magnesium decreased (36%), edema (34%), nausea (33%), dyspnea (32%), uric acid increased (32%), potassium decreased (32%), alkaline phosphatase increased (30%), mucositis (28%), aspartate aminotransferase increased (27%), phosphatase decreased (25%), electrocardiogram QT prolonged (24%), rash (24%), creatinine increased (24%), cough (23%), decreased appetite (22%), myalgia (21%), constipation (20%), and pyrexia (20%).
- **In patients with newly diagnosed AML**, the most frequently reported Grade ≥3 adverse reactions (≥5%) were fatigue (14%), differentiation syndrome (11%), electrocardiogram QT prolonged (11%), diarrhea (7%), nausea (7%), and leukocytosis (7%). Serious adverse reactions (≥5%) were differentiation syndrome (18%), electrocardiogram QT prolonged (7%), and fatigue (7%). There was one case of posterior reversible encephalopathy syndrome (PRES).
- **In patients with relapsed or refractory AML**, the most frequently reported Grade ≥3 adverse reactions (≥5%) were differentiation syndrome (13%), electrocardiogram QT prolonged (10%), dyspnea (9%), leukocytosis (8%), and tumor lysis syndrome (6%). Serious adverse reactions (≥5%) were differentiation syndrome (10%), leukocytosis (10%), and electrocardiogram QT prolonged (7%). There was one case of progressive multifocal leukoencephalopathy (PML).

### DRUG INTERACTIONS

**Strong or Moderate CYP3A4 Inhibitors:** Reduce TIBSOVO® dose with strong CYP3A4 inhibitors. Monitor patients for increased risk of QTc interval prolongation.

**Strong CYP3A4 Inducers:** Avoid concomitant use with TIBSOVO®.

**Sensitive CYP3A4 Substrates:** Avoid concomitant use with TIBSOVO®.

**QTc Prolonging Drugs:** Avoid concomitant use with TIBSOVO®. If co-administration is unavoidable, monitor patients for increased risk of QTc interval prolongation.

## LACTATION

Because many drugs are excreted in human milk and because of the potential for adverse reactions in breastfed children, advise women not to breastfeed during treatment with TIBSOVO® and for at least 1 month after the last dose.

**Please see full Prescribing Information, including Boxed WARNING.**

## About Agios

Agios is focused on discovering and developing novel investigational medicines to treat malignant hematology, solid tumors and rare genetic diseases through scientific leadership in the field of cellular metabolism. In addition to an active research and discovery pipeline across these three therapeutic areas, Agios has two approved oncology precision medicines and multiple first-in-class investigational therapies in clinical and/or preclinical development. For more information, please visit the company's website at [www.agios.com](http://www.agios.com).

## Cautionary Note Regarding Forward-Looking Statements

This press release contains forward-looking statements within the meaning of The Private Securities Litigation Reform Act of 1995. Such forward-looking statements include those regarding: the potential benefits of TIBSOVO® (ivosidenib); Agios' plans to submit a supplemental new drug application for TIBSOVO® in previously treated cholangiocarcinoma between the end of 2020 and mid-2021; and Agios' strategic plans and prospects. The words "anticipate," "believe," "estimate," "expect," "intend," "may," "plan," "predict," "project," "would," "could," "potential," "possible," "hope" and similar expressions are intended to identify forward-looking statements, although not all forward-looking statements contain these identifying words. Such statements are subject to numerous important factors, risks and uncertainties that may cause actual events or results to differ materially from Agios' current expectations and beliefs. For example, there can be no guarantee that any product candidate Agios or its collaborators is developing will successfully commence or complete necessary preclinical and clinical development phases, or that development of any of Agios' product candidates will successfully continue. There can be no guarantee that any positive developments in Agios' business will result in stock price appreciation. Management's expectations and, therefore, any forward-looking statements in this press release could also be affected by risks and uncertainties relating to a number of other important factors, including, without limitation: risks and uncertainties related to the impact of the COVID-19 pandemic to Agios' business, operations, strategy, goals and anticipated milestones, including its ongoing and planned research activities, ability to conduct ongoing and planned clinical trials, clinical supply of current or future drug candidates, commercial supply of current or future approved products, and launching, marketing and selling current or future approved products; Agios' results of clinical trials and preclinical studies, including subsequent analysis of existing data and new data received from ongoing and future studies; the content and timing of decisions made by the U.S. FDA, the EMA or other regulatory authorities, investigational review boards at clinical trial sites and publication review bodies; Agios' ability to obtain and maintain requisite regulatory approvals and to enroll patients in its planned clinical trials; unplanned cash requirements and expenditures; competitive factors; Agios' ability to obtain, maintain and enforce patent and other intellectual property protection for any product candidates it is developing; Agios' ability to maintain key collaborations; and general economic and market conditions. These and other risks are described in greater detail under the caption "Risk Factors" included in Agios' public filings with the Securities and Exchange Commission. Any forward-looking statements contained in this press release speak only as of the date hereof, and Agios expressly disclaims any obligation to update any forward-looking statements, whether as a result of new information, future events or otherwise, except as required by law.

## Contacts

### Investors:

Holly Manning, 617-844-6630  
Director, Investor Relations  
[Holly.Manning@agios.com](mailto:Holly.Manning@agios.com)

### Media:

Jessica Rennekamp, 857-209-3286  
Associate Director, Corporate Communications  
[Jessica.Rennekamp@agios.com](mailto:Jessica.Rennekamp@agios.com)



Source: Agios Pharmaceuticals, Inc.