

Cautionary Statements

This presentation and various remarks we make during this presentation contain forward-looking statements within the meaning of The Private Securities Litigation Reform Act of 1995. Such forward-looking statements include those regarding Agios' plans, strategies and expectations for its and its collaborator's preclinical, clinical and commercial advancement of its drug development programs including TIBSOVO® (ivosidenib), IDHIFA® (enasidenib), mitapivat, vorasidenib, AG-270, and AG-946; the potential benefits of Agios' product candidates; its key milestones and guidance for 2020; its strategic vision and goals for 2025; its plans regarding future data presentations; its financial guidance regarding the period in which it will have capital available to fund its operations; and the potential benefits of its strategic plans and focus. The words "anticipate," "expect," "goal," "hope," "milestone," "opportunity," "plan," "potential," "possible," "strategy," "will," "vision," and similar expressions are intended to identify forward-looking statements, although not all forward-looking statements contain these identifying words. Such statements are subject to numerous important factors, risks and uncertainties that may cause actual events or results to differ materially from Agios' current expectations and beliefs. For example, there can be no guarantee that any product candidate Agios or its collaborators is developing will successfully commence or complete necessary preclinical and clinical development phases, or that development of any of Agios' product candidates will successfully continue. There can be no guarantee that any positive developments in Agios' business will result in stock price appreciation. Management's expectations and, therefore, any forward-looking statements in this presentation and various remarks we make during this presentation could also be affected by risks and uncertainties relating to a number of other important factors, including, without limitation: risks and uncertainties related to the impact of the COVID-19 pandemic to Agios' business, operations, strategy, goals and anticipated milestones, including its ongoing and planned research activities, ability to conduct ongoing and planned clinical trials, clinical supply of current or future drug candidates, commercial supply of current or future approved products, and launching, marketing and selling current or future approved products; Agios' results of clinical trials and preclinical studies, including subsequent analysis of existing data and new data received from ongoing and future studies; the content and timing of decisions made by the U.S. FDA, the EMA or other regulatory authorities, investigational review boards at clinical trial sites and publication review bodies; Agios' ability to obtain and maintain requisite regulatory approvals and to enroll patients in its planned clinical trials; unplanned cash requirements and expenditures; competitive factors; Agios' ability to obtain, maintain and enforce patent and other intellectual property protection for any product candidates it is developing; Agios' ability to maintain key collaborations; and general economic and market conditions. These and other risks are described in greater detail under the caption "Risk Factors" included in Agios' public filings with the SEC. Any forward-looking statements contained in this presentation and various remarks we make during this presentation speak only as of the date hereof, and Agios expressly disclaims any obligation to update any forward-looking statements, whether as a result of new information, future events or otherwise, except as required by law.



We are driven by our sense of urgency to help patients.







On a bad day, it's like watching some electronic toy slowly lose the battery.

—Tamara S., Minnesota




The disease has affected my career. I spent 11 years to get a PhD in nutrition...My heart wants more but my body can't handle it.

—Tamara S., Minnesota

Currently 50 years old. Diagnosed with PK deficiency at the age of 6.


Our Strategy is Clear

For more than a decade, our mission has been to create differentiated, small molecule medicines for patients in three focus areas – malignant hematology, solid tumors and rare genetic diseases – based on our unique expertise in cellular metabolism and adjacent areas of biology





Our People and Culture Fuel Incredible Productivity, Strategic Focus and Continuity from Early Research to Market





Agios 2025 Vision: Focused Innovation. Ambitious Development. Transformative Treatments for Patients Across Three Focus Areas.

NOW 2025 **COMMERCIAL MEDICINES MEDICINES** 8+ LABEL **EXPANSION INDICATIONS INDICATIONS PRODUCTIVE** 6+ **DISCOVERY MOLECULES IN THE CLINIC ENGINE MOLECULES IN THE CLINIC** \$105-115M **CASH FLOW FINANCIAL EXPECTED U.S. TIBSOVO® POSITIVE 2020 REVENUE**



- Finalize pivotal development plan for mitapivat in thalassemia by the end of 2020; initiate a pivotal program in 2021
- Finalize pivotal development plan for mitapivat in sickle cell disease and initiate pivotal program in 2021
- Initiate Phase 1 study of AG-946, next-generation PKR activator, in healthy volunteers in mid-2020
- Mitapivat ACTIVATE and ACTIVATE-T topline data in PK deficiency between YE 2020 and mid-2021

- Achieve full-year U.S. revenue for TIBSOVO® \$105-115M
- EU regulatory process for TIBSOVO® in R/R AML ongoing with a CHMP opinion expected by YE 2020


SOLID TUMORS


Mature OS data from ClarIDHy study of TIBSOVO® in mid-2020; cholangio sNDA filing between YE 2020 and mid-2021

Anticipated


Milestones and

Recent Updates






Significant Growth Potential in Malignant Hematology




TIBSOVO®

R/R AML	U.S. Approval; MAA Under Review
1L Monotherapy	U.S. Approval
1L HMA Combo	Phase 3
1L 7+3 Combo	Phase 3





Significant Q1 Growth Driven by Increased Demand in Both R/R and Frontline AML Segments and Expanding Customer Base





16% Growth

In Product Revenue Quarter-over-Quarter



\$105 - 115M

U.S. Net Sales Guidance for 2020



25% Increase

In Unique Prescribers Quarter-over-Quarter



~1,400


Patients Treated Since Launch


Source: Agios estimates



Advancing
Toward Largest
Opportunity for
mIDH1 AML:
Intensive and
Non-Intensive
Therapy
Combinations

50K AML Patients Diagnosed Per Year in U.S. and EU





Four Distinct Solid Tumor Opportunities Across Three Clinical Molecules



TIBSOVO®


R/R Cholangio

sNDA expected YE 2020 to mid-2021



Low-grade Glioma

Phase 3



AG-270

Phase 1

Combo

2nd Line NSCLC




AG-270

1st or 2nd Line Phase 1 Pancreatic Cancer Combo



Established Utility of IDH Inhibition in Solid Tumors with Positive ClarIDHy Phase 3 Study of TIBSOVO® in Second-line or Later Cholangiocarcinoma

Mature OS from ClarIDHy Phase 3 expected mid-2020; sNDA planned between YE 2020 and mid-2021





Global Phase 3 INDIGO Study of Vorasidenib in IDH Mutant Low-Grade Glioma

SIGNIFICANT 2-HG SUPPRESSION IMPRESSIVE PRELIMINARY EFFICACY DATA

ENCOURAGING PHASE 1 DATA

>90%


2-HG
SUPPRESSION IN
RESECTED
mIDH1 GLIOMAS
ACROSS ALL
DOSES TESTED

33% ORR

IN THE
VORASIDENIB
ARM OF THE
PERIOPERATIVE
STUDY

22 mo.

MEDIAN TREATMENT DURATION IN VORASIDENIB PHASE 1




Endpoints

Primary: Progression free survival (by BIRC)

Secondary/Exploratory: Tumor volume, safety, ORR, OS, QOL, seizures, neuro-cognitive function, time to next intervention



AG-270, MAT2A Inhibitor, Preclinical Data Supports Combination with Taxanes; Two Phase 1 Combination Arms Ongoing




PHASE 1 COMBINATION ARMS INITIATED


AG-270 + docetaxel in MTAP-deleted NSCLC (2nd line) N = up to 40


AG-270 + nab-paclitaxel and gemcitabine in MTAP-deleted pancreatic ductal adenocarcinoma (1st or 2nd line) N = up to 45






PKR Activation Represents Unique Mechanism of Action with Potential to Address Broad Range of Hemolytic Anemias





Inadequate Production of ATP

 PKR mutations decrease PK stability, ATP generation and RBC membrane integrity and increase RBC destruction, leading to chronic hemolytic anemia




Increased Demand of ATP

 In other hemolytic anemias, there is an increase in ATP demand and impaired ATP production, leading to damage and premature death of RBCs, hemolysis and anemia



Mitapivat has Potential to be First Disease-modifying Therapy for Patients with PK Deficiency





Chronic daily dosing with mitapivat for a median of 3 years and up to 42 months was well tolerated

COMPLICATIONS & COMORBIDITIES REGARDLESS OF TRANSFUSION STATUS

SUPPORTIVE CARE ONLY

HIGH RISK OF IRON OVERLOAD


HIGHER LIFETIME
RATES OF
PULMONARY
HYPERTENSION,
OSTEOPOROSIS,
AND LIVER
CIRRHOSIS

O APPROVED THERAPIES 38%
OF PATIENTS NOT RECEIVING REGULAR
TRANSFUSIONS EXPERIENCE IRON OVERLOAD

Source: Data presented at ASH 2019; van Beers EJ, et al. Haematologica. 2019;104(2):e51-e53.



PKR Activators May Improve Thalassemic RBC Production and Survival by Increasing ATP Production





Study Design: Open-label, Phase 2, Multi-center Trial of Mitapivat in Thalassemia

Key Inclusion Criteria


- β-thalassemia ± α-globin gene mutations, HbE β-thalassemia, or αthalassemia (HbH disease)
- Hb \leq 10.0 g/dL
- Non–transfusion-dependent^a

Primary Endpoint^b

 Hb response, defined as increase of ≥ 1.0 g/dL from baseline at any time between weeks 4–12, inclusive

Secondary/Exploratory Endpoints


 Sustained Hb response; delayed Hb response; markers of hemolysis; hematopoietic activity; safety





Demographics and Disposition

Baseline characteristics	Total (N = 18)
Median (range) duration of treatment, weeks	20.6 (1.1–50.0)
Male/female, n	5/13
Age at informed consent, median (range), years	43.5 (29–67)
Race, n (%) Asian White Native Hawaiian or other Pacific Islander Other ^a	9 (50.0) 4 (22.2) 1 (5.6) 4 (22.2)
Thalassemia type, n (%) α β	5 (27.8) 13 (72.2)
Hb baseline, median (range), g/dL	8.43 (5.6–9.8)
Indirect bilirubin, median (range), mg/dL	1.17 (0.31–5.52)
Lactate dehydrogenase, median (range), U/L	249 (126–513)
Erythropoietin, median (range), mU/mL	70.5 (15–11,191)



Interim Phase 2 Results in Thalassemia: Primary Endpoint Met in 92.3% of Patients

Endpoint	Genotype	N/N	%	90% CI
Ub responders during weeks 4, 12 among these who completed 12	All	12/13	92.3	68.4, 99.6
Hb responders during weeks 4–12 among those who completed 12 weeks	α	4/4	100	47.3, 100
	β	8/9	88.9	57.1, 99.4
Hb responders during weeks 12–24 among those who completed 24 weeks	βª	8/9	88.9	57.1, 99.4
Sustained responders: primary response and ≥ 2 Hb responses during weeks 12–24	βª	7/8	87.5	52.9, 99.4

Patient population	N	Weeks	Mean (SD) change from baseline Hb, g/dL
All patients	13	4–12	1.34 (0.7)
α-thalassemia	4	4–12	1.17 (0.4)
β-thalassemia	9	4–24	1.43 (0.8)
β-thalassemia responders	8	4-24	1.63 (0.5)
All responders	12	4-12	1.47 (0.5)

Interim Phase 2 Results in Thalassemia: Activation of wPKR by Mitapivat Improved Hb and Associated Markers of Hemolysis and Erythropoiesis

Treatment with mitapivat induced Hb increase of ≥1.0 g/dL in 92% evaluable patients, including 100% α-thalassemia patients; 7 of 8 evaluable patients achieved sustained Hb response

Median (range) time to Hb increase of ≥ 1 g/dL among responders was 3.1 (1.4–7.1) weeks


Mitapivat was generally well tolerated; the safety profile was consistent with previous studies

Improvements in markers of hemolysis and erythropoiesis correlated with the Hb increases

Mean ATP percent increase from baseline was similar to that previously observed with mitapivat in healthy volunteers



PKR Activation in Sickle Cell Disease Modulates 2,3-DPG and ATP to Potentially Improve Anemia and Reduce Sickling





NHLBI & Agios CRADA Study of Mitapivat in SCD: Study Design

Primary

Safety and tolerability

- Frequency and severity of adverse events
- Changes in laboratory parameters (including reticulocyte counts and levels of hemoglobin, bilirubin, and lactate dehydrogenase)


Secondary


Pharmacokinetics/pharmacodynamics

- Pharmacokinetics of mitapivat (AG-348)
- Levels of 2,3-DPG, PK-R, and ATP, and oxygen dissociation sickling in RBCs
- Relationship between mitapivat pharmacokinetics and safety

Safety, Tolerability, Pharmacokinetics, and Pharmacodynamics of Escalating Multiple Oral Doses of Mitapivat in Subjects With Stable SCD

ClinicalTrials.gov NCT04000165: Nonrandomized, open-label, phase 1 study; N ≈ 15–25







Clinical Proof-of-concept for Mitapivat Established in Sickle Cell Disease

7 of 8 (88%) efficacy evaluable patients experienced a Hb increase, and 5 of 8 (63%) patients achieved a Hb increase of ≥1.0 g/dL from baseline (range 1.0-2.7 g/dL) at doses of 50 mg BID or lower.


Treatment with mitapivat was associated with decreases in hemolytic markers such as bilirubin, LDH and reticulocytes.


2,3-DPG decreases and increases in ATP levels were observed. Sickling curves (t50) and oxygen dissociation curves (p50) consistent with decreases in both sickling and HbS polymerization.


AEs generally consistent with previously reported data with mitapivat treatment or are to be expected in the context of SCD. One SAE, a VOC, occurred during drug taper and was possibly attributed to mitapivat.




Illustrative Sickle Cell Patient Case Study: Male, 39 Years Old












PKR Activation Has Potential Broad Utility Across Hemolytic Anemias








NTD β- and α- Thalassemia	Phase 2 enrollment complete
Thalassemia	Pivotal plan expected by YE and initiation in 2021



Adult SCD	NIH CRADA; data to be submitted to ASH
Adult SCD	Pivotal study expected to initiate in 2021





First Quarter 2020 Financial Results

Statement of Operations	Three Months Ended 3/31/20	Three Months Ended 3/31/19	Year Ended 12/31/19
Total Revenue	\$87.1M	\$30.2M	\$117.9M
Collaboration Revenue TIBSOVO® Net Sales Royalty Revenue	61.1M 22.7M 3.3M	18.9M 9.1M 2.2M	47.5M 59.9M 10.5M
Cost of Sales	0.5M	0.3M	1.3M
Research & Development Expense	91.3M	95.6M	410.9M
Selling, General & Administrative Expense	38.5M	31.8M	132.0M

Balance Sheet	3/31/20	12/31/19
Cash, Cash Equivalents and Marketable Securities	\$613.1M	\$717.8M



agios

AGIOS 2025 VISION:

Focused Innovation. Ambitious Development.

Transformative Treatments for Patients Across Three Focus Areas.

